



COOKED MOLASSES TUB RECEIVING TUB WITH AMAFERM®

For Stressed Cattle

GUARANTEED ANALYSIS:

Crude Protein, Min	10.0%
Crude Fat, Min	5.0%
Crude Fiber, Max	2.5%
Calcium (Ca), Min	2.0%
Calcium (Ca), Max	2.5%
Phosphorus (P), Min	1.0%
Salt (NaCl), Min.....	2.0%
Salt (NaCl), Max.....	2.5%
Magnesium (Mg), Min	1.0%
Potassium (K),Min	2.5%
Cobalt (Co), Min	145 ppm
Copper (Cu), Min.....	1,000 ppm
Iodine (I), Min	40 ppm
Manganese (Mn), Min	2,000 ppm
Selenium (Se), Min	13.2 ppm
Zinc (Zn), Min	2,900 ppm
Vitamin A, Min	200,000 IU/lb
Vitamin D ₃ Min	40,000 IU/lb
Vitamin E, Min	800 IU/lb

CAUTION:

Consumption of this product by sheep and goats may result in copper toxicity. Feeding selenium at levels in excess of 0.3 ppm in the total diet is prohibited. Use only as directed.



Amaferm® is a registered trademark of Biozyme, Inc.

INGREDIENTS:

Molasses Products, Processed Grain By-Products, Monocalcium Phosphate, Saccharomyces cerevisiae Yeast Culture, Calcium Carbonate, Salt, Magnesium Oxide, Dried Aspergillus oryzae Fermentation Product, Zinc Amino Acid Complex, Manganese Amino Acid Complex, Copper Amino Acid Complex, Zinc Sulfate, Manganese Sulfate, Basic Copper Chloride, Vitamin E Supplement, Vitamin A Supplement, Vitamin D₃ Supplement, Niacin Supplement, Calcium Pantothenate, Biotin, Riboflavin Supplement, Pyridoxine Hydrochloride, Thiamine Mononitrate, Folic Acid, Vitamin B12 Supplement, Selenium Yeast, Ethylenediamine Dihydroiodide, Cobalt Glucoheptonate, Cobalt Carbonate, Lactic Acid and Diatomaceous Earth.

FEEDING DIRECTIONS:

Feed RANCHER'S CHOICE® Receiving Cooked Molasses Tub with Amaferm® free-choice to stressed cattle. Make sure animals have adequate pasture or roughage available and are not starved for salt or minerals. Feed continuously. Recommended consumption of RANCHER'S CHOICE® Receiving Cooked Molasses Tub with Amaferm® is 6 to 8 ounces per head per day. Consumption will vary according to the weight of the animal and the quality and amount of available forage. Increase the number of tubs and move tubs closer to water or loafing areas to increase intake. Decrease the number of tubs or move farther from water or loafing areas to decrease intake. Provide at least one tub for every 15 to 20 animals. Provide free access to fresh, clean water at all times.